

Download

Literaturliste zum Beitrag: „Ohne Worte verständigen“

Astrid Steinmetz, *pfllegen: palliativ* 39: 8 ff



- Alsawy, S.; Mansell, W.; McEvoy, P.; Tai, S. (2017): *What is good communication for people living with dementia? A mixed-methods systematic review*. *International Psychogeriatrics*, 29(11): 1785–1800
- Asplund, K.; Norberg, A.; Adolfsson, R.; Waxman, H. M. (1991): *Facial expressions in severely demented patients: A stimulus-response study of four patients with dementia of Alzheimer type*. *International Journal of Geriatric Psychiatry*, 6: 599–606
- Barnes, S.; Gardiner, C.; Gott, M.; Payne, S.; Chady, B.; Small, N.; Seamark, D.; Halpin, D. (2012): *Enhancing patient-professional communication about end-of-life issues in life-limiting conditions: A critical review of the literature*. *Journal of Pain and Symptom Management*, 44 (6): 866–879
- Bradshaw, J. (2001): *Complexity of staff communication and reported level of understanding skills in adults with intellectual disability*. *Journal of Intellectual Disability Research*, 45: 233–243
- Cull, A.; Hay, C.; Love, S. B.; Mackie, M.; Smets, E.; Stewart, M. (1996): *What do cancer patients mean when they complain of concentration and memory problems?* *British Journal of Cancer*, 74 (10): 1674–1679
- Decety, J.; Lamm, C. (2009): *The Biological Bases of Empathy*. In: Cacioppo, J. T.; Berntson, G. G. (Eds.): *Handbook of Neuroscience for the Behavioral Sciences*. New York: John Wiley and Sons: 940–957
- Eritz, H.; Hadjistavropoulos, T. (2011): *Do informal caregivers consider nonverbal behavior when they assess pain in people with severe dementia?* *The Journal of Pain*, 12 (3): 331–339
- Fuchs, T. (2003): *Non-verbale Kommunikation: Phänomenologische, entwicklungspsy-chologische und therapeutische Aspekte*. *Z. klin. Psychol. Psychiat. Psychother.* 51: 333–345
- Gottschlich, M. (2007). *Medizin und Mitgefühl: Die heilsame Kraft empathischer Kommunikation*. (2., vollst. überarb. Auflage). Wien: Böhlau
- Gutknecht, D. (2010): *Professionelle Responsivität: Ein hochschulbezogenes Ausbildungskonzept für den frühpädagogischen Arbeitskontext U3: Kinder unter drei Jahren und ihre Familien*. Pädagogische Hochschule Heidelberg. Heidelberg. Unter: <http://d-nb.info/1008380415/34> (eingesehen am 19.03.2018)
- Healy, D.; Noonan Walsh, P. (2007): *Communication among nurses and adults with severe and profound intellectual disabilities: Predicted and observed strategies*. *Journal of Intellectual Disabilities*, 11 (2): 127–141

- Johnson, A.; Chang, E.; Daly, J.; Harrison, K.; Noel, M.; Hancock, K.; Easterbrook, S. K.; Easterbrook, S. (2009): *The communication challenges faced in adopting a palliative care approach in advanced dementia*. International Journal of Nursing Practice, 15: 467–474
- Kessler, H.; Bayerl, P.; Deighton, R. M.; Traue, H. C. (2002): *Facially Expressed Emotion Labeling (FEEL): PC-gestützter Test zur Emotionserkennung*. In: Verhaltenstherapie und Verhaltensmedizin, 23(3): 297–306
- Kruse, A. (1992): *Sterbende begleiten: Anthropologische Überlegungen, psychologische Beiträge und Erarbeitung von psychologischen Grundlagen einer Sterbebegleitung*. In: Schmitz-Scherzer, R. (Hrsg.). Altern und Sterben: 63–103. Bern: Hans Huber
- Mayer, J. D.; Salovey, P.; Caruso, D. R.; Sitarenios, G. (2003): *Measuring emotional intelligence with the MSCEIT V2.0*. Emotion, 3 (1): 97–105
- Moore, P. M.; Rivera Mercado, S.; Grez Artigues, M.; Lawrie, T.A. (2013): *Communication skills training for healthcare professionals working with people who have cancer*. Cochrane Database of Systematic Reviews, 3, Art. No.: CD003751. DOI: 10.1002/14651858.CD003751.pub3. [23.06.2018]
- Ng, K.; Gunten, C. F. von. (1998): *Symptoms and attitudes of 100 consecutive patients admitted to an acute hospice/palliative care unit*. Journal of Pain and Symptom Management, 16 (5): 307–316
- Pappas, J. B.; Pappas, E. C. (2015): *The Sustainable Personality: Values and Behaviors in Individual Sustainability*. International Journal of Higher Education, 4(1): 12–21
- Riggio, R. E. (2006): *Nonverbal Skills and Abilities*. In: Patterson, M. L.; Manusov, V. L. (Hrsg.): The SAGE Handbook of Nonverbal Communication: 79–95. Thousand Oaks, Calif., London: Sage
- Roter, D. L.; Frankel, R. M.; Hall, J. A.; Sluyter, D. (2006): *The expression of emotion through nonverbal behavior in medical visits: Mechanisms and outcomes*. Journal of General Internal Medicine, 21: 28–34
- Schofield, N. G.; Green, C.; Creed, F. (2008): *Communication skills of health care professionals working in oncology – Can they be improved?* European Journal of Oncology Nursing, 12: 4–13
- Schore, J.R.; Schore, A.N. (2008): *Modern attachment theory: the central role of affect regulation in development and treatment*.
- Soothill, K.; Morris, S. M.; Harman, J.; Francis, B.; Thomas, C.; McIlmurray, M. B. (2001): *The significant unmet needs of cancer patients: Probing psychosocial concerns*. Support Cancer Care, 9: 597–605
- Steinmetz, A. (2016): *Kommunikation ohne Worte – KoW®. Nonverbale Interaktion mit demenzkranken und palliativen Patienten*. Wiesbaden: Springer VS
- Stern, D.N. (1998): *Die Lebenserfahrungen des Säuglings*. 6. Aufl. Stuttgart: Klett-Cotta

- Suchman, A. L.; Markakis, K.; Beckman, H. B.; Frankel, R. (1997): *A model of empathic communication in the medical interview*. The Journal of the American Medical Association, 277 (8): 678–682
- Tuffrey-Wijne, I.; McEnhill, L. (2008): *Communication difficulties and intellectual disability in end-of-life care*. International Journal of Palliative Nursing, 14 (4): 189–194
- Voltz, R.; Borasio, G. D. (2001): *Neuropsychiatrische Symptome in der Palliativmedizin*. Schmerz, 15 (5): 339–343
- Wagener, R.; Berkemeyer, C.; Hock, G.; Schneider, A.; Winterberg, C.; Ulmer, E.M. (1998): *Essen und Trinken bei Menschen mit Alzheimer-Demenz. Eine Interaktionsstudie*. Pflege, 11 (2): 89–95
- Zimmermann, C.; Del Piccolo, L.; Finset, A. (2007): *Cues and concerns by patients in medical consultations: A literature review*. Psychological Bulletin, 133 (3): 438–463